

Peri-operative management of anticoagulation and antiplatelet therapy

David Keeling,¹ R. Campbell Tait,² and Henry Watson³ on behalf of the British Committee for Standards in Haematology

¹Oxford University Hospitals NHS Foundation Trust, Oxford, ²Glasgow Royal Infirmary, Glasgow, and ³Aberdeen Royal Infirmary, Aberdeen, UK

Keywords: perioperative, anticoagulation, antiplatelet, warfarin, surgery.

Cardiovascular Society; these organisations do not necessarily approve or endorse the contents.

Methodology

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) nomenclature was used to evaluate levels of evidence and to assess the strength of recommendations. The GRADE criteria are specified in the British Committee for Standards in Haematology (BCSH) guidance pack (http://www.bcshguidances.com/BCSH_PROCESS/42_EVIDENCE_LEVELS_AND_GRADES_OF_RECOMMENDATION.html) and the GRADE working group website <http://www.gradeworkinggroup.org>.

Literature review details

Details of the literature review are given in Appendix 1.

Review of manuscript

Review of the manuscript was performed by the British Society for Haematology (BSH) Guidelines Haemostasis and Thrombosis Task Force, BSH Guidelines Executive Committee and by the Haemostasis and Thrombosis sounding board of the BSH. The latter comprises 50 or more members of the BSH who have commented on the content and applicability in the UK setting. It has also been sent to the following organisations for review: The Royal College of Surgeons; The Royal College of Anaesthetists; Thrombosis UK, a patient-centred charity dedicated to promoting awareness, research & care of thrombosis; the British Dental Association; Intra-Health (who operate National Health Service General Practitioner and community pharmacies); and the British

Introduction

A BSH guideline on warfarin (Keeling *et al*, 2011) addressed the issue of perioperative management and is updated in this article to include the issue of perioperative management of patients on direct oral anticoagulants (DOACs) and antiplatelet agents, which are becoming frequent clinical queries. This guideline will consider whether and when anticoagulants and antiplatelet agents should be stopped before elective surgery and invasive procedures, when agents can be restarted and how to manage patients on these drugs who require emergency surgery. If an anticoagulant or antiplatelet effect persists, haemostasis may be improved by the use of pre-operative parenteral tranexamic acid, which has been shown to reduce blood loss and transfusion requirements in both cardiac and trauma surgery, without increasing thrombotic complications (McIlroy *et al*, 2009; Shakur *et al*, 2010).

For agents with a slow offset and onset of action, bridging therapy with an alternative drug at a full treatment dose can be considered in patients deemed to be at high risk of thrombosis; this mainly concerns whether treatment dose low molecular weight heparin (LMWH) or unfractionated heparin (UFH) should be given when warfarin is temporarily discontinued. Thromboprophylaxis with low dose LMWH is not regarded as 'bridging'.

For some invasive procedures, such as dentistry (Perry *et al*, 2007) (see also <http://www.sdcep.org.uk/published-guidance/anticoagulants-and-antiplatelets/>), joint injections (Ahmed & Gertner, 2011), cataracts (Jamula *et al*, 2009), pacemaker insertion (Ahmed *et al*, 2010; Airaksinen *et al*, 2013) and certain endoscopic procedures (Veitch *et al*, 2016), anticoagulation may not need to be stopped. Procedures that require anticoagulation to be stopped will vary in their bleeding risk and, importantly, the consequences of bleeding will depend on the site of surgery and local anatomy. Although some have grouped procedures into lower or higher risk (Spyropoulos & Douketis, 2012; Baron *et al*, 2013) we think the operating surgeon, dentist, or interventional radiologist has to assess the risk of bleeding for the

Correspondence: BCSH Secretary, British Society for Haematology, 100 White Lion Street, London, N1 9PF, UK.
E-mail: bcsh@b-s-h.org.uk

individual patient and discuss both this and the plan for peri-operative anticoagulation with them. The plan must be recorded clearly in the notes, including a plan for when the patient is discharged.

Warfarin and other vitamin K antagonists

Warfarin has a half-life of approximately 36 h and as its effect subsides γ -carboxylated vitamin K-dependent procoagulant factors need to be synthesised. Warfarin therefore needs to be stopped 5 days before elective surgery to ensure haemostasis has returned to normal. This is likely to differ for other vitamin K antagonists with different half-lives (acenocoumarol, 10 h; phenindione, 8 h; fluindione, 3 days; phenprocoumon, 5 days). If possible, the International Normalized Ratio (INR) should be determined the day before surgery to allow the administration of phytomenadione if the INR is ≥ 1.5 , so reducing the risk of cancellation. The INR should be checked on the day of surgery. Stopping warfarin for a shorter time and attempting to reverse its effect with oral phytomenadione on the day before surgery did not prove a satisfactory alternative (Steib *et al*, 2010). Due to its slow onset of action warfarin can be resumed, at the normal maintenance dose (Douketis *et al*, 2012), or with two initial days of double maintenance dose (Schulman *et al*, 2014), the evening of surgery (or the next day) if there is adequate haemostasis.

There have been many reviews and attempts to estimate the risk of peri-operative thrombosis (Dunn & Turpie, 2003; Dunn *et al*, 2007; Dentali *et al*, 2012; Douketis *et al*, 2012; Siegal *et al*, 2012; Spyropoulos & Douketis, 2012). The main question has been whether the risk of thrombosis is sufficiently high, in patients who have temporarily discontinued a coumarin, to use treatment dose LMWH or UFH pre-operatively and/or post-operatively when haemostasis is secure. This is predicated on the assumption that bridging will reduce the thrombotic risk. It is noteworthy that in their meta-analysis Siegal *et al* (2012) found no difference in the risk of thromboembolic events in eight studies comparing bridged and non-bridged groups of patients (odds ratio [OR], 0.80; 95% confidence interval [CI], 0.42–1.54). However, bridging was associated with an increased risk of major bleeding in five studies (OR, 3.60; 95% CI, 1.52–8.50). A further systematic review also concluded 'while the antithrombotic efficacy of perioperative bridging with LMWH has not been demonstrated, increased bleeding risk is observed in different types of surgery' (Eijgenraam *et al*, 2013).

Patients taking warfarin for the treatment and secondary prevention of venous thromboembolism (VTE), for stroke prevention in atrial fibrillation (AF) or for mechanical heart valves (MHV) need separate consideration. For patients with acute VTE the risk of recurrence without anticoagulation is very high in the first 3 months (Kearon & Hirsh, 1997) and surgery will increase the risk further. When a patient is more than 3 months from an acute event the risk of recurrence is

much lower, the treatment phase is over, and patients are remaining on an anticoagulant for secondary prevention (Kearon & Akl, 2014). Prophylactic dose LMWH can substitute for warfarin after the acute treatment period, hence patients with VTE more than 3 months prior can usually simply be given post-operative prophylactic dose LMWH (or a suitable alternative) rather than receive full dose bridging therapy while anticoagulation with a coumarin is re-established. Bridging might be considered for those thought to be at very high risk, such as patients with a previous VTE occurring whilst on therapeutic anticoagulation who now have a target INR of 3–5.

For patients with a MHV the risk varies with type of valve (bileaflet less than caged ball and tilting disc), valve position (aortic less than mitral) and patient risk factors (such as previous stroke or transient ischaemic attack (TIA), AF and reduced left ventricular ejection fraction). We have previously recommended bridging therapy for patients with MHVs other than those with a bileaflet aortic valve and no other risk factors (Keeling *et al*, 2011) and there is no strong evidence to change this recommendation.

For patients with atrial fibrillation the CHADS₂ (Congestive failure; Hypertension; Age ≥ 75 years; Diabetes mellitus; prior Stroke, TIA or thromboembolism) score or, more recently, the CHA₂DS₂-VASc CHADS₂ + Vascular disease; Age 65–74 years; Sex category) score has been used to predict stroke risk. The CHADS₂ score may also predict risk of post operative stroke (Kaatz *et al*, 2011) and guidelines have suggested the CHADS₂ score is used to select patients for bridging (Douketis *et al*, 2012) whilst the previous BCSH guideline suggested bridging in only those with a previous stroke or TIA or multiple other risk factors (Keeling *et al*, 2011). There is now a randomized, double-blind, placebo-controlled trial of bridging in AF patients (Douketis *et al*, 2015). Patients were randomised to dalteparin (100 iu/kg bd) or placebo from 3 days before until 24 h before the procedure and then for 5–10 days after the procedure. A total of 1884 patients were enrolled, with 950 assigned to receive no bridging therapy and 934 assigned to receive bridging therapy. The incidence of arterial thromboembolism was 0.4% in the no-bridging group and 0.3% in the bridging group (risk difference, 0.1 percentage points; 95% CI –0.6 to 0.8; $P = 0.01$ for non-inferiority). The incidence of major bleeding was 1.3% in the no-bridging group and 3.2% in the bridging group (relative risk, 0.41; 95% CI 0.20–0.78; $P = 0.005$ for superiority). The authors concluded forgoing bridging was non-inferior to bridging for the prevention of arterial thromboembolism and decreased the risk of major bleeding. Thirty-eight per cent of patients had a CHADS₂ score of ≥ 3 , though only 3.1% had a score ≥ 5 (which requires a previous stroke or TIA plus at least three of four other risk factors), 9.4% had a previous stroke and 8.3% a previous TIA. Patients with a stroke or TIA within the previous 12 weeks were excluded. Our updated advice is shown in Table I. When we say 'consider bridging' we do not mean it

Guideline

Table I. When to consider bridging with treatment dose heparin in patients who stop warfarin if thrombotic risk is especially high.

	Consider bridging with treatment dose heparin in
VTE	Patients with a VTE within previous 3 months. Very high risk patients such as patients with a previous VTE whilst on therapeutic anticoagulation who now have a target INR of 3–5.
AF	Patients with a previous stroke/TIA in last 3 months. Patients with a previous stroke/TIA and three or more of the following risk factors: <ul style="list-style-type: none"> • Congestive cardiac failure • Hypertension (>140/90 mmHg or on medication) • Age >75 years • Diabetes mellitus
MHV	MHV patients other than those with a bileaflet aortic valve and no other risk factors

VTE, venous thromboembolism; INR, International Normalized Ratio; AF, atrial fibrillation; TIA, transient ischaemic attack; MHV, mechanical heart valve.

should be automatically given, but consider whether to bridge or not in discussion with the patient.

In patients who are receiving pre-operative bridging with LMWH, the last dose should be at least 24 h before surgery and if on a once a day regimen, some recommend the last dose is halved for high risk surgery (Douketis *et al*, 2012). We recommend that post-operative bridging (i.e. full dose anticoagulation) is not started until at least 48 h after high bleeding risk surgery although thromboprophylaxis should be given if indicated.

Emergency surgery in patients on warfarin

If surgery can wait for 6–8 h then 5 mg of intravenous phytonadione can restore coagulation factors; if this is not possible, anticoagulation can be reversed with 25–50 u/kg of four-factor prothrombin complex concentrate (Refaai *et al*, 2013; Goldstein *et al*, 2015), we would give at the lower end of this range and check the INR.

Post-operative management should follow the same strategy as for elective surgery.

Recommendations

- **Warfarin should be stopped for 5 days before an elective procedure if anticoagulation needs to be discontinued (1C).**
- **Patients with venous thromboembolism (VTE) more than 3 months earlier can usually be given post-operative prophylactic dose low molecular weight heparin (LMWH) (or a suitable alternative) rather than bridging therapy (2C).**
- **Patients at very high risk of recurrent VTE, such as patients with a previous VTE whilst on therapeutic**

anticoagulation who now have a target International Normalized Ratio (INR) of 3–5, and patients who have had VTE less than 3 months previously should be considered for bridging (2D)

- **Patients with atrial fibrillation who have a CHADS₂ score of ≤4 and who have not had a stroke or transient ischaemic attack (TIA) in last three months should not receive bridging (1A)**
- **Patients with a bileaflet aortic mechanical heart valve (MHV) with no other risk factors do not require bridging whilst it should be considered in all other MHV patients (2C).**
- **We recommend that post-operative bridging is not started until at least 48 h after high bleeding risk surgery (1C).**

Direct oral anticoagulants

The approach to the peri-operative management of patients on Direct oral anticoagulants (DOACs) is based on an approximate calculation of the half-life of the drug and therefore its persistence in the circulation, taking into account renal function. This is combined with consideration of the bleeding risk of the proposed procedure and a clinical evaluation of the patient's individual risk factors for thrombosis and bleeding. Current strategies for elective surgery do not routinely include measurement of either non-specific or specific coagulation parameters to assist in quantification of DOAC levels. For each of the drugs there are data on periods of discontinuation during the studies to evaluate the drug against coumarin therapy. In addition there is a single moderate sized, prospective study evaluating the outcomes of a set protocol for the peri-procedural management of patients on dabigatran (Schulman *et al*, 2015).

Dabigatran

During the Randomized Evaluation of Long-Term Anticoagulation Therapy (RE-LY) study treatments were transiently discontinued in 4591 subjects (Healey *et al*, 2012), i.e. around 25% of the study population. A range of procedures was performed – mostly cardiac catheterisation, dental extraction and colonoscopy but also more major surgery. The incidence of major bleeding was 3.8%, 5.1% and 4.6% respectively for dabigatran 110 mg, dabigatran 150 mg and warfarin and the rate of stroke and systemic embolism was 0.5% in each group for the 30-day period around the discontinuation. Significant bleeding and thrombosis were more common after emergency and major procedures.

A multicentre prospective study that included 324 standard risk and 217 high risk procedures and which broadly followed the protocol in Table II reported low rates of major bleeding, 1.8% (0.7–3%) and thromboembolism, 0.2% (0–0.5%) in the 30-day period around the procedure

(Schulman *et al*, 2015). Dabigatran was restarted post-operatively only when haemostasis had been secured. After minor procedures it recommenced at 75 mg on the evening of the procedure escalating to 110 or 150 mg BD the following day. For major procedures the re-introduction was again guided by haemostasis, with most patients restarting at normal full dose 48–72 h post-procedure. Bridging with LMWH or UFH was used in 1.7% of cases post-procedure but not at all pre-procedure. In this study, 40% of the procedures had a high bleeding risk and eight of the 10 patients who had major bleeding had undergone a high bleeding risk procedure.

Rivaroxaban

Data from the Rocket-AF study show that in 2130 patients undergoing 2980 procedures, periods of discontinuation of rivaroxaban for ≥ 3 days to allow surgery or an invasive procedure result in no significant difference in any of, major haemorrhage, clinically relevant non-major haemorrhage, stroke and systemic embolism, myocardial infarction and death when compared to discontinuation of warfarin (Sherwood *et al*, 2014). There are no published data that report on a fixed protocol for peri-procedural discontinuation of rivaroxaban but several groups have proposed schedules for this (Lai *et al*, 2014; Heidbuchel *et al*, 2015). The manufacturer's advice is to discontinue rivaroxaban for over 24 and 48 h respectively for low bleeding risk and high bleeding risk procedures.

Apixaban

Data from the Aristotle trial also described 9260 procedures where anticoagulation may be interrupted (Garcia *et al*, 2014). The vast majority of the events were low risk procedures, and apixaban was discontinued for around 60% of these. Again the major outcomes were similar in patients taking warfarin and apixaban irrespective of whether or not anticoagulation was discontinued. Of note in this observational study, when apixaban was discontinued it was for 2–5 days. There are no published data that report on a fixed protocol for peri-procedural discontinuation of apixaban. The manufacturer's advice is to discontinue apixaban for over 24 and 48 h respectively for low bleeding risk and high bleeding risk procedures. Published schedules on the discontinuation of apixaban give different advice. Lai *et al* (2014) and Heidbuchel *et al* (2015) take renal function into consideration and suggest minimum periods of discontinuation for those with creatinine clearance (CrCl) of 15–30 ml/min undergoing high risk procedures of over 48 h while Ward *et al* (2013) did not consider renal function but recommend discontinuation for high bleeding risk procedures for at least 60 h.

In making recommendations we have considered the practicalities of instructing patients around periods of discontinuation. While Lai *et al* (2014) and the European Heart Rhythm Association guideline (Heidbuchel *et al*, 2015)

suggest a period of discontinuation for 36 h in patients with CrCl of 15–30 ml/min undergoing low risk procedures, this is not really practical for patients taking a once daily preparation, such as rivaroxaban, and so we have chosen to recommend a 48-h period of discontinuation for this group when using either apixaban or rivaroxaban.

Edoxaban

The most recent oral Xa inhibitor to be licenced in the UK is edoxaban, 50% of which is renally excreted and its half-life is 10–14 h. The Summary of Product Characteristics suggests that for surgical or other procedures it should be discontinued preferably at least 24 h before the procedure.

Emergency surgery in patients on DOACs

There are few data on the management of emergency surgery in patients receiving DOACs. The ability to make predictions regarding haemostasis at surgery in these patients is limited, firstly, by uncertainty in the concentration of each drug that is associated with haemostatic safety. Secondly, a UK National External Quality Assessment Service (NEQAS) supplementary exercise undertaken in October 2014 found high coefficients of variation for the Haemoclot assay and for the chromogenic anti-Xa assays when assaying plasma levels of DOACs (personal communication, Dr S Kitchen, UK NEQAS, Sheffield, UK). The greatest variation was seen in the measurement of low concentrations of drug, and, worryingly, drug was reported as being present in samples where there was no drug present. If an anticoagulant effect cannot be excluded neuroaxial anaesthesia should be avoided.

When possible, surgery should be delayed to allow the plasma level of the drug to fall. The concentration of drug can be estimated from the dose of the drug, time of last dose and the patients' renal function. Other factors, such as patient sex, weight and the use of interacting drugs, will also have less significant effects. The approximate half-life of the drugs according to renal function is given in Table II.

Coagulation tests. Examination of routine coagulation tests such as the prothrombin time (PT), activated partial thromboplastin time (APTT) and thrombin time (TT) may allow an approximate estimate of the levels of drug present in the circulation (Kitchen *et al*, 2014). A normal TT effectively excludes the presence of dabigatran in a sample but normal APTT and PT do not exclude the presence of significant concentrations of rivaroxaban, apixaban or edoxaban in a sample. Guidance would then suggest that advanced or non-routine assays can be applied to the situation but with caution given to the interpretation of the results in view of the caveats mentioned above (Kitchen *et al*, 2014).

Prohaemostatic agents. It has been suggested that use of several pro-haemostatic agents might reduce the risk of peri-

Table II. Usual time to discontinue DOACs before surgery or invasive procedures for which anticoagulation needs to be stopped.

Renal function (CrCl, ml/min)	Estimated half-life (h)	Low bleeding risk (h)	High bleeding risk (h)
Dabigatran			
≥80	13	24	48
≥50 to <80	15	24–48	48–72
≥30 to <50	18	48–72	96
Rivaroxaban			
≥30	9	24	48
<30		48	72
Apixaban			
≥30	8	24	48
<30		48	72
Edoxaban			
≥30	10–14	24	48
<30		48	72

DOAC, direct oral anticoagulant; CrCl, creatinine clearance.

surgical bleeding in patients on DOACs who require emergency surgery. Most of the evidence cited in this regard relates to animal experiments and to the observation of changes in *in vitro* tests of haemostasis. The usual view of clinicians is that treatment with a prothrombin complex concentrate (PCC) might improve outcomes, but this doesn't take into account the potential for adverse thrombotic outcomes, which are often overlooked. There are few data that strongly support the use of PCC and activated PCC in the management of emergency surgery (Makris, 2014) and so a pragmatic approach might be to proceed with surgery considering PCC only in the event of diffuse coagulopathic bleeding. Tranexamic acid is likely to reduce bleeding and should be given.

Other strategies. Other general management strategies include avoiding further intake of anticoagulants and avoiding the use of any additional therapies such as nonsteroidal anti-inflammatory drugs (NSAID) and colloids (dextrans and starches) that might further compromise haemostasis.

Dabigatran is minimally protein bound and so can be removed by dialysis if a procedure can be delayed for long enough for this to take place, but this is rarely practical. Significant amounts of dabigatran can be removed by a single dialysis session although rebound increases in concentration have been observed on cessation of dialysis (Chang *et al*, 2013; Chai-Adisaksotha *et al*, 2015). This strategy is not applicable to rivaroxaban, apixaban and edoxaban, which are all highly protein bound.

Specific reversal agents. *Idarucizumab*—In a prospective study of the administration of a standard dose of 5 g of idarucizumab to 36 patients who were receiving dabigatran prior to undergoing an invasive or surgical procedure, normal intraoperative haemostasis was reported in 33, and

mildly or moderately abnormal haemostasis was reported in 2 patients and 1 patient, respectively. One thrombotic event occurred within 72 h after idarucizumab. In the same study, major reversal of dabigatran effect on coagulation tests was observed in patients given idarucizumab in 88–98% of patients (Pollack *et al*, 2015).

Andexanet—There are no data on the use of andexanet in patients undergoing surgery but there are now promising data on the reversal of anticoagulation in healthy volunteers. In a study of apixaban- and rivaroxaban-treated volunteers who received a bolus dose of andexanet, significant reductions in anticoagulant activity were seen in both groups (Siegal *et al*, 2015). In apixaban-treated individuals, anti-factor Xa activity was reduced by 94% compared with 21% among those who received placebo and unbound apixaban concentration fell by 9.3 µg/l compared with 1.9 µg/l. Thrombin generation was fully restored in 100% of andexanet recipients and 11% of placebo recipients. In rivaroxaban-treated volunteers, anti-factor Xa activity was reduced by 92% compared with 18% among those who received placebo, and unbound rivaroxaban concentration fell by 23.4 µg/l compared with 4.2 µg/l. Thrombin generation was fully restored in 96% of andexanet and 7% of placebo recipients. These effects were sustained when andexanet was administered as a bolus plus an infusion. No serious adverse or thrombotic events were witnessed. Although these data cannot be directly interpreted as being able to secure haemostasis during surgery, the findings are promising (Siegal *et al*, 2015).

Recommendations

- Patients with normal renal function undergoing planned low risk procedures should not take a direct oral anticoagulant (DOAC) for 24 h before the procedure (2B)
- Patients with normal renal function undergoing planned higher risk procedures should not take a DOAC for 48 h before the procedure (2B)
- For patients with renal impairment see Table II (2D).
- Following minor or low risk procedures in patients with low bleeding risk, anticoagulation can be recommenced 6–12 h post-procedure if haemostasis has been fully secured (2C)
- Following high risk procedures and in patients with an increased bleeding risk or in any situation where any increased risk of bleeding is unacceptable, DOACs should not be re-introduced at full dose until at least 48 h post-procedure (2C)
- In patients with high thrombosis risk it is appropriate to consider prophylactic doses of anticoagulation before re-introducing full therapeutic dose DOAC (2D)
- DOAC measurement by indirect methods using dilute thrombin time, ecarin clotting time and calibrated anti-Xa assays should currently be interpreted with caution

in the management of patients receiving a DOAC who require emergency surgery (2B)

- A normal thrombin time can be interpreted as indicating that there is a minimal circulating concentration of dabigatran. Normal prothrombin time (PT) and activated partial thromboplastin time (APTT) do not exclude significant concentrations of dabigatran, rivaroxaban or apixaban (1A)
- If an anticoagulant effect cannot be excluded, neuroaxial anaesthesia should be avoided (1C).
- Prothrombin complex concentrates should not be routinely used in patients on DOACs prior to emergency surgery (2D)
- Tranexamic acid is likely to reduce bleeding in patients who have a residual anticoagulant effect (1C).
- Drugs and colloids that impair the haemostatic mechanism should be avoided in the peri-surgical management of patients receiving DOACs (2D)
- Idarucizumab should be used to reverse dabigatran therapy prior to emergency invasive procedures and surgery where the bleeding risk is considered significant (1C)
- Andexanet, when available, should be used to reverse apixaban, rivaroxaban or edoxaban prior to emergency invasive procedures and surgery where the bleeding risk is considered significant (2C)

Antiplatelet therapy

Antiplatelet therapy is a key pharmacological intervention in the secondary prevention of cardiovascular disease. This pertains particularly to clopidogrel following ischaemic cerebrovascular disease and dual antiplatelet therapy (DAPT) following acute coronary syndromes (ACS) when a combination of aspirin and an ADP receptor (P2Y₁₂) antagonist is indicated, especially after coronary artery stenting. In these situations the small day-to-day increase in bleeding risk associated with aspirin, and more so with DAPT (Sorensen *et al*, 2009), is outweighed by their clinical benefit. However, the continuation of antiplatelet agents in the surgical setting is associated with an increase in bleeding risk. In a meta-analysis of 41 studies Burger *et al* (2005) demonstrated that aspirin therapy was associated with a 1.5-fold increase in post-operative bleeding events, but no increase in the severity of bleeds – concluding that low dose aspirin could be continued through most surgical procedures except neurosurgery and prostatectomy. A recent meta-analysis has shown that patients on clopidogrel who have a hip fracture can be managed by normal protocols with early surgery (Soo *et al*, 2016). However, the same may not be true for DAPT, which is associated with significantly more surgery-related bleeding (14.7%) compared to aspirin (4.1%) (Singla *et al*, 2012). This bleeding risk has to be balanced against the considerable increased thrombotic risk associated with premature termination or interruption of anti-platelet monotherapy (Biondi-

Zoccai *et al*, 2006) or DAPT (Mehran *et al*, 2013; Rossini *et al*, 2015), which may be required to facilitate a surgical or other invasive procedure.

Non-cardiac surgical procedures

Three randomized controlled trials (RCTs) in high-risk elective surgery have compared temporary peri-operative interruption or continuation of aspirin in patients with stable cardiovascular disease, very few participants had experienced recent (<30 days) ACS or had undergone a stenting procedure (Oscarsson *et al*, 2010; Mantz *et al*, 2011; Devereaux *et al*, 2014). Two of the studies were terminated early with small numbers recruited and were therefore underpowered to assess differences in bleeding events (Oscarsson *et al*, 2010; Mantz *et al*, 2011). Omitting aspirin from 10 days prior to surgery until the morning of surgery resulted in no increase in thrombotic events or decrease in bleeding events (Mantz *et al*, 2011). In contrast, omitting aspirin from 7 days prior to surgery until 3 days post-op resulted in a significantly higher 30-day rate of Major Adverse Cardiac Events (MACE) [9% vs. 1.8%, $P = 0.02$] but no difference in peri-operative blood loss (Oscarsson *et al*, 2010). Consensus views in guidelines have generally recommend continuation of aspirin monotherapy unless surgery is perceived to have a particularly high bleeding risk or is in a confined space such as brain, posterior eye chamber or medullary canal (Korte *et al*, 2011). More recently a larger RCT including 4382 patients already on an antiplatelet agent demonstrated no difference in the composite endpoint of death or non-fatal myocardial infarction (hazard ratio [HR] 1.00 (0.81–1.23)) nor in major bleeding (HR 1.11 (0.84–1.48) if aspirin was continued as opposed to being withheld from 1 day pre-op until 7 days post-op (Devereaux *et al*, 2014). Major bleeding was increased in a separate group not on aspirin but randomized to start it (HR 1.34 (1.03–1.74). There is very limited and less reliable data on bleeding risks and cardiovascular benefits of continuing single agent clopidogrel peri-operatively (Luckie *et al*, 2009). While some guidelines propose continuing clopidogrel monotherapy in the same situations as with aspirin monotherapy (Ferraris *et al*, 2012), we do not believe there is sufficient evidence to make a recommendation.

More challenging is the management of DAPT around invasive procedures. This is an increasingly common scenario as more patients undergo percutaneous intervention (PCI) with stent insertion following ACS, when DAPT is recommended for at least 4 weeks following bare metal stent and 12 months following drug-eluting stent (although shorter duration DAPT is required with the newer bioabsorbable polymer drug-eluting stents). Between 4% and 8% of PCI patients will require surgery within 1 year of stenting. The risk of peri-operative MACE is greatest within the first month after PCI with gradually lessening risk at 2–6 months, 6–12 months and >1 year (Nuttall *et al*, 2008; Savonitto *et al*, 2011). Other recognized markers for post-op MACE

include interruption of antiplatelet therapy, recent ACS, urgent or high cardiac-risk surgery and chronic kidney disease (Albaladejo *et al*, 2011; Rossini *et al*, 2015). There are no RCTs on which to base advice in this setting, hence most guidelines adopt a pragmatic expert consensus view, using a matrix assessing the patient's thrombotic risk and the bleeding risk associated with the type of invasive procedure being undertaken (Korte *et al*, 2011; Rossini *et al*, 2014).

In summary, very low bleeding-risk procedures can be undertaken without stopping DAPT, whereas low risk procedures in patients with low thrombotic risk may be undertaken on aspirin with temporary cessation of the ADP receptor antagonist. Ideally, elective surgery in patients deemed to be at high thrombotic risk should be deferred until they are lower risk. If surgery cannot be deferred then it should generally proceed on aspirin with temporary discontinuation of the ADP receptor antagonist. In high thrombotic risk patients requiring high bleeding-risk surgery that cannot be deferred, consideration can be given to bridging with a parenteral short-acting glycoprotein IIb/IIIa inhibitor, such as tirofiban or eptifibatide, during the period of ADP receptor antagonist withdrawal (Savonitto *et al*, 2011).

Cardiovascular surgery

The cardiovascular benefit of aspirin following coronary artery bypass graft (CABG) surgery is well established. In a meta-analysis of 13 studies the pre-op use of aspirin was associated with a significant reduction in ischaemic events (OR 0.56, 0.33–0.96) but with small increases in post-op bleeding and transfusion requirements and a 1.85-fold increased risk of re-exploration surgery (Hastings *et al*, 2015). In contrast, in a recent randomized trial, the administration of preoperative aspirin to patients undergoing coronary artery surgery did not result in a lower risk of death or thrombotic complications or in a higher risk of bleeding as compared to placebo (Myles *et al*, 2016). Continuing clopidogrel pre-CABG was also shown, in a meta-analysis of 11 cohort studies, to increase post-op chest tube drainage and the requirement for blood products as well as 2- to 5-fold increase in re-exploration rates (Kunadian *et al*, 2006). In both these meta-analyses the study groups were heterogeneous and poorly controlled. Indeed, in the clopidogrel analyses it is understood that many of the cases were also on aspirin. Furthermore, many of the included studies were pre-2000 when use of tranexamic acid, which has been shown to reduce blood loss in both aspirin- and clopidogrel-treated patients (McIlroy *et al*, 2009; Shi *et al*, 2013), was less prevalent. Hence, most recent cardiac guidelines recommend continuing aspirin pre-CABG unless there is a very high bleeding risk, a very low thrombosis risk or the patient would decline transfusion (Ferraris *et al*, 2012; Sousa-Uva *et al*, 2014).

Cardiac surgery under combined aspirin and clopidogrel is complicated by a further increase in bleeding, compared to

aspirin alone. Compared to aspirin and clopidogrel DAPT, aspirin and ticagrelor DAPT had similar CABG-related bleeding rates while aspirin and prasugrel DAPT was associated with higher bleeding and surgical re-exploration rates (Held *et al*, 2011; Smith *et al*, 2012). Therefore, based on observational data on bleeding and drug metabolite half lives, it is recommended that clopidogrel and ticagrelor are discontinued 5 days pre-op and prasugrel 7 days pre-op while aspirin is continued throughout (Ferraris *et al*, 2012; Capodanno & Angiolillo, 2013; Sousa-Uva *et al*, 2014). In particularly high thrombotic risk patients, bridging protocols with a short-acting parenteral antiplatelet agent during withdrawal of the oral ADP receptor antagonist have been proposed (Savonitto *et al*, 2011; Capodanno & Angiolillo, 2013). The parenteral glycoprotein IIb/IIIa inhibitor is usually commenced on day –3 and stopped 4–6 h pre-op, and commenced 4–6 h post-op until the ADP receptor antagonist can be restarted (within 7 days, when all bleeding has been controlled), and in the case of clopidogrel an initial loading dose is recommended (Sousa-Uva *et al*, 2014).

Emergency surgery in patients on antiplatelet therapy

When urgent high haemorrhage-risk surgery is indicated, and time does not permit cessation of one or both antiplatelet agents, there is evidence from both small *in vitro* (Vilahrur *et al*, 2007; Li *et al*, 2012) and *in vivo* (Thiele *et al*, 2012) studies that transfusion of donor platelets may improve haemostasis. Platelets should be transfused at least 2 h after the last dose of aspirin and 12–24 h after the last dose of clopidogrel to avoid being inhibited by circulating drug or active metabolite. Reversal of aspirin requires fewer donor platelets and is more complete because aspirin-inactivated platelets can be recruited by thromboxane generated in transfused platelets, so whilst 2 pools of platelets reverse the effect of aspirin, the effect of even higher doses of platelets when on ADP antagonists is less certain (Vilahrur *et al*, 2007; Li *et al*, 2012; Hansson *et al*, 2014; Godier *et al*, 2015). A single dose of platelets in patients with an intracranial haemorrhage on aspirin did not improve outcome (Baharoglu *et al*, 2016).

Inter-individual variation in sensitivity to ADP receptor antagonists, particularly clopidogrel, may allow some patients to have a shorter period of discontinuation pre-operatively (e.g. 3 days for clopidogrel or ticagrelor and 5 days for prasugrel). This is particularly relevant in urgent surgery situations when assessment of platelet function may help identify those patients in which early surgery may be safer. Corredor *et al* (2015) have reviewed the utility of pre-op platelet function testing. A great variety of devices are available, however the most robust clinical evidence appears to be with thromboelastography platelet mapping or multiple electrode aggregometry. With the latter, a normal pre-op platelet aggregation response to ADP was associated with low post-op bleeding (Ranucci *et al*, 2011). Using thromboelastography platelet mapping, platelet receptor inhibition (PRI) of

>76% was associated with an 11-fold higher risk of transfusion while a PRI <34% had a negative predictive value of 90% for requiring at least 2 units red cell transfusion (Kwak *et al*, 2010; Kasivisvanathan *et al*, 2014).

Neuroaxial anaesthesia

Horlocker *et al* (2010), on behalf of the American Society of Regional Anesthesia, reviewed several large studies of neuroaxial anaesthesia in a variety of surgical and obstetric settings. Although spinal haematoma has been reported following spinal or epidural anaesthesia, such events are rare – estimated incidence <1 in 150 000 epidural and <1 in 220 000 spinal anaesthetics. In a series of 61 cases of spinal haematoma, antiplatelet therapy was only implicated in three and furthermore, no cases of spinal haematoma were reported in combined series of >6000 patients having central neural blockade while on antiplatelet therapy. Hence the guideline recommends spinal and epidural anaesthesia can be undertaken without cessation of NSAID or aspirin (Horlocker *et al*, 2010). In a review of practice this was done for low-risk (e.g. peripheral nerve blocks) but not for high-risk pain management procedures (Narouze *et al*, 2015). Limited evidence is currently available in relation to the safety of neuro-axial anaesthesia in patients receiving ADP-receptor antagonists, and therefore it is recommended that all such agents be discontinued 7 days prior to the procedure (Horlocker *et al*, 2010; Narouze *et al*, 2015).

Recommendations

- **When being used for secondary prevention of cardiovascular disease, aspirin monotherapy can be continued for most invasive non-cardiac procedures (including neuroaxial anaesthesia) but, if the perceived bleeding risk is high, aspirin can be omitted from day –3 to day +7 with no net detriment (2C)**
- **Aspirin can be continued both before and after coronary artery bypass surgery (1B)**
- **Hip fracture surgery can take place early in patients on clopidogrel (1B)**
- **For urgent low bleeding risk surgery in patients on antiplatelet agents routine platelet transfusion should not be given (2C)**
- **For urgent high-bleeding risk surgery in patients on antiplatelet agents**
 - **Given the uncertain net benefit of platelet transfusion, consider the use of pre-operative intravenous tranexamic acid (2C)**
 - **If, despite tranexamic acid, there is excessive peri- or post-op bleeding, or if the bleeding risk is perceived to be very high, consider infusion of 2 pools of donor platelets. This may improve haemostasis if given at least two h after the last dose of aspirin though even**

higher doses of donor platelets 12–24 h after the last dose of clopidogrel may have a lesser effect (2C)

- **In patients with a recent acute coronary syndrome or coronary artery stent on dual antiplatelet therapy low bleeding risk procedures should proceed without interruption of antiplatelet therapy (1C)**
- **In patients with a recent acute coronary syndrome or coronary artery stent on dual antiplatelet therapy elective high bleeding risk procedures should, if possible, be postponed in patients still requiring dual antiplatelet therapy (1C). If surgery cannot be deferred, aspirin should be continued and clopidogrel or ticagrelor interrupted from 5 days pre-op or prasugrel from 7 days pre-op (1C).**

Disclaimer

While the advice and information in this guidance is believed to be true and accurate at the time of going to press, neither the authors, the BSH, nor the publishers accept any legal responsibility for the content of this guidance.

Acknowledgments

All the authors reviewed the literature and wrote the initial draft of the manuscript. The authors wish to thank Jackie Wilson for help in undertaking the initial literature review. The BSH Haemostasis and Thrombosis task force members at the time of writing this guidance were D Perry, M Laffan, E Chalmers, A Mumford, I Jennings, K Gomez, R Alikhan, W. Lester, I Walker and E Gray. The authors would like to thank them and the BSH sounding board, BSH Guidelines executive committee for their support in preparing this guideline.

Declaration of interests

The BSH Guidelines paid the expenses incurred during the writing of this guidance (see http://www.bcsbguidelines.com/BCSH_PROCESS/DOCUMENTS_FOR_TASK_FORCE-S_AND_WRITING_GROUPS/203_Expense_forms_and_policy.html). DK has been paid for attending advisory boards by Bayer, Boehringer Ingelheim, Pfizer, Daiichi Sanko, Octapharma and CSL Behring. RCT has been paid for attending advisory boards by Bayer, Boehringer Ingelheim and BMS/Pfizer.

Review process

Members of the writing group will inform the writing group Chair if any new pertinent evidence becomes available that would alter the strength of the recommendations made in this document or render it obsolete. The document will be archived and removed from the BSH current guidelines website if it becomes obsolete. If new recommendations are

Guideline

made an addendum will be published on the BSH guidelines website.

Audit tool

See http://www.bcsbguidelines.com/4_HAEMATOLOGY_GUIDELINES.html for template

Appendix 1

Search overview for perioperative management of anticoagulation and antiplatelet therapy

MEDLINE

1. exp Platelet Aggregation Inhibitors/
2. (antiplatelet therapy or Aspirin or Dipyridamole or Ticlopidine or Clopidogrel or Prasugrel or Ticagrelor).tw.
3. (platelet* adj1 (antagonist* or inhibitor* or antiaggregant*)).tw.
4. (antiplatelet adj1 (drug* or agent*)).tw.
5. exp Anticoagulants/
6. (anticoagula* or 4-hydroxycoumarins or Acenocoumarol or Dicumarol or Heparin or Heparin, low-molecular-weight or Phenindione or Phenprocoumon or Warfarin or Dabigatran or Rivaroxaban or Apixaban or Edoxaban).tw.
7. (thrombin* adj2 inhibitor*).tw.
8. or/1–7
9. exp Perioperative Care/
10. (perioperative or periprocedural).tw.
11. (pre-operative or intra-operative or post-operative or pre operative or intra operative or post operative).tw.
12. or/9–11
13. 8 and 12
14. (surgery or operation* or procedure* or intervention* or treatment*).tw.
15. exp specialties, surgical/
16. 14 or 15
17. 13 and 16
18. (animals not (humans and animals)).sh.
19. 17 not 18
20. limit 19 to english language

EMBASE

1. exp antithrombocytic agent/
2. (antiplatelet therapy or Aspirin or Dipyridamole or Ticlopidine or Clopidogrel or Prasugrel or Ticagrelor).tw.
3. (platelet* adj1 (antagonist* or inhibitor* or antiaggregant*)).tw.
4. (antiplatelet adj1 (drug* or agent*)).tw.

5. exp anticoagulant agent/
6. (anticoagula* or 4-hydroxycoumarins or Acenocoumarol or Dicumarol or Heparin or Heparin, low-molecular-weight or Phenindione or Phenprocoumon or Warfarin or Dabigatran or Rivaroxaban or Apixaban or Edoxaban).tw.
7. (thrombin* adj2 inhibitor*).tw.
8. or/1–7
9. exp perioperative period/
10. (perioperative or periprocedural).tw.
11. (pre-operative or intra-operative or post-operative or pre operative or intra operative or post operative).tw.
12. or/9–11
13. 8 and 12
14. (surgery or operation* or procedure* or intervention* or treatment*).tw.
15. exp specialties, surgical/
16. 14 or 15
17. 13 and 16
18. (animal/or nonhuman/) not human/
19. 17 not 18
20. limit 19 to english language
21. limit 20 to embase

Records generated

Database searched	Date searched	Results
MEDLINE (OVID) 1946	16/4/15	4349
EMBASE (OVID) 1974–2015 March 31	16/4/15	9145
Total		13494
After de-duplication		11084
Further duplicates removed during review (136)		10948
Exclusions (9693)		1255
Non-English (1)		
Non-human (15)		
Paediatric (5)		
Not relevant to clinical question (9580)		
Too few cases (92)		

Breakdown of remaining results

Articles included	1003
Antiplatelet only (380)	
New oral anticoagulants (NOAC) only (69)	
Vitamin K antagonists (VKA) only (271)	
Both NOAC and VKA (63)	
Mixed antiplatelet and anticoagulant (220)	
Possible articles	252
Possibly relevant based on title only (159)	
Surveys of practice (77)	
Relating to service or costs (16)	

References

- Ahmed, I. & Gertner, E. (2011) Safety of arthrocentesis in patients on chronic warfarin therapy with therapeutic INR. *Journal of Hospital Medicine*, **2**, S1.
- Ahmed, I., Gertner, E., Nelson, W.B., House, C.M., Dahiya, R., Anderson, C.P., Benditt, D.G. & Zhu, D.W.X. (2010) Continuing warfarin therapy is superior to interrupting warfarin with or without bridging anticoagulation therapy in patients undergoing pacemaker and defibrillator implantation. *Heart Rhythm: the Official Journal of the Heart Rhythm Society*, **7**, 745–749.
- Airaksinen, K.E.J., Korkeila, P., Lund, J., Ylitalo, A., Karjalainen, P., Virtanen, V., Raatikainen, P., Koivisto, U.-M. & Koistinen, J. (2013) Safety of pacemaker and implantable cardioverter-defibrillator implantation during uninterrupted warfarin treatment—the FinPAC study. *International Journal of Cardiology*, **168**, 3679–3682.
- Albaladejo, P., Marret, E., Samama, C.-M., Collet, J.-P., Abhay, K., Loutrel, O., Charbonneau, H., Jaber, S., Thoret, S., Bosson, J.-L. & Piriou, V. (2011) Non-cardiac surgery in patients with coronary stents: the RECO study. *Heart*, **97**, 1566–1572.
- Baharoglu, M.I., Cordonnier, C., Salman, R.A., de Gans, K., Koopman, M.M., Brand, A., Majoie, C.B., Beenen, L.F., Marquering, H.A., Vermeulen, M., Nederkoorn, P.J., de Haan, R.J. & Roos, Y.B. (2016) Platelet transfusion versus standard care after acute stroke due to spontaneous cerebral haemorrhage associated with antiplatelet therapy (PATCH): a randomised, open-label, phase 3 trial. *Lancet*, **387**, 2605–2613.
- Baron, T.H., Kamath, P.S. & McBane, R.D. (2013) Management of antithrombotic therapy in patients undergoing invasive procedures. *New England Journal of Medicine*, **368**, 2113–2124.
- Biondi-Zoccai, G.G., Lotrionte, M., Agostoni, P., Abbate, A., Fusaro, M., Burzotta, F., Testa, L., Sheiban, I. & Sangiorgi, G. (2006) A systematic review and meta-analysis on the hazards of discontinuing or not adhering to aspirin among 50,279 patients at risk for coronary artery disease. *European Heart Journal*, **27**, 2667–2674.
- Burger, W., Chemnitz, J.M., Kneissl, G.D. & Rucker, G. (2005) Low-dose aspirin for secondary cardiovascular prevention - Cardiovascular risks after its perioperative withdrawal versus bleeding risks with its continuation - Review and meta-analysis. *Journal of Internal Medicine*, **257**, 399–414.
- Capodanno, D. & Angiolillo, D.J. (2013) Management of antiplatelet therapy in patients with coronary artery disease requiring cardiac and noncardiac surgery. *Circulation*, **128**, 2785–2798.
- Chai-Adisaksoha, C., Hillis, C., Lim, W., Boonyawat, K., Moffat, K. & Crowther, M. (2015) Hemodialysis for the treatment of dabigatran-associated bleeding: a case report and systematic review. *Journal of Thrombosis and Haemostasis*, **13**, 1790–1798.
- Chang, D.N., Dager, W.E. & Chin, A.I. (2013) Removal of dabigatran by hemodialysis. *American Journal of Kidney Diseases*, **61**, 487–489.
- Corredor, C., Wasowicz, M., Karkouti, K. & Sharma, V. (2015) The role of point-of-care platelet function testing in predicting postoperative bleeding following cardiac surgery: a systematic review and meta-analysis. *Anaesthesia*, **70**, 715–731.
- Dentali, F., Pignatelli, P., Malato, A., Poli, D., Di Minno, M.N., Di Gennaro, L., Rancan, E., Pastori, D., Grifoni, E., Squizzato, A., Siragusa, S., Di Minno, G. & Ageno, W. (2012) Incidence of thromboembolic complications in patients with atrial fibrillation or mechanical heart valves with a subtherapeutic international normalized ratio: a prospective multicenter cohort study. *American Journal of Hematology*, **87**, 384–387.
- Devereaux, P.J., Mrkobrada, M., Sessler, D.I., Leslie, K., Alonso-Coello, P., Kurz, A., Villar, J.C., Sigamani, A., Biccard, B.M., Meyhoff, C.S., Parlow, J.L., Guyatt, G., Robinson, A., Garg, A.X., Rodseth, R.N., Botto, F., Lurati Buse, G., Xavier, D., Chan, M.T.V., Tiboni, M., Cook, D., Kumar, P.A., Forget, P., Malaga, G., Fleischmann, E., Amir, M., Eikelboom, J., Mizera, R., Torres, D., Wang, C.Y., VanHelder, T., Paniagua, P., Berwanger, O., Srinathan, S., Graham, M., Pasin, L., Le Manach, Y., Gao, P., Pogue, J., Whitlock, R., Lamy, A., Kearon, C., Baigent, C., Chow, C., Pettit, S., Chrolavicius, S., Yusuf, S. & Investigators, P. (2014) Aspirin in patients undergoing noncardiac surgery. *New England Journal of Medicine*, **370**, 1494–1503.
- Douketis, J.D., Spyropoulos, A.C., Spencer, F.A., Mayr, M., Jaffer, A.K., Eckman, M.H., Dunn, A.S. & Kunz, R. (2012) Perioperative management of antithrombotic therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*, **141**, e326S–e350S.
- Douketis, J.D., Spyropoulos, A.C., Kaatz, S., Becker, R.C., Caprini, J.A., Dunn, A.S., Garcia, D.A., Jacobson, A., Jaffer, A.K., Kong, D.F., Schulman, S., Turpie, A.G., Hasselblad, V. & Ortel, T.L. (2015) Perioperative bridging anticoagulation in patients with atrial fibrillation. *New England Journal of Medicine*, **373**, 823–833.
- Dunn, A.S. & Turpie, A.G. (2003) Perioperative management of patients receiving oral anticoagulants: a systematic review. *Archives of Internal Medicine*, **163**, 901–908.
- Dunn, A.S., Spyropoulos, A.C. & Turpie, A.G. (2007) Bridging therapy in patients on long-term oral anticoagulants who require surgery: the Prospective Peri-operative Enoxaparin Cohort Trial (PROSPECT). *Journal of Thrombosis and Haemostasis*, **5**, 2211–2218.
- Eijgenraam, P., Ten Cate, H. & Ten Cate-Hoek, A. (2013) Safety and efficacy of bridging with low molecular weight heparins: a systematic review. *Journal of Thrombosis and Haemostasis*, **11**, 13.
- Ferraris, V.A., Saha, S.P., Oestreich, J.H., Song, H.K., Rosengart, T., Reece, T.B., Mazer, C.D., Bridges, C.R., Despotis, G.J., Joiner, K. & Clough, E.R. (2012) 2012 update to the Society of Thoracic Surgeons guideline on use of antiplatelet drugs in patients having cardiac and noncardiac operations. *Annals of Thoracic Surgery*, **94**, 1761–1781.
- Garcia, D., Alexander, J.H., Wallentin, L., Wojdyla, D.M., Thomas, L., Hanna, M., Al-Khatib, S.M., Dorian, P., Ansell, J., Commerford, P., Flaker, G., Lanus, F., Vinereanu, D., Xavier, D., Hylek, E.M., Held, C., Verheugt, F.W.A., Granger, C.B. & Lopes, R.D. (2014) Management and clinical outcomes in patients treated with apixaban vs warfarin undergoing procedures. *Blood*, **124**, 3692–3698.
- Godier, A., Taylor, G. & Gaussem, P. (2015) Inefficacy of platelet transfusion to reverse ticagrelor. *New England Journal of Medicine*, **372**, 196–197.
- Goldstein, J.N., Refaai, M.A., Milling, T.J. Jr, Lewis, B., Goldberg-Alberts, R., Hug, B.A. & Sarode, R. (2015) Four-factor prothrombin complex concentrate versus plasma for rapid vitamin K antagonist reversal in patients needing urgent surgical or invasive interventions: a phase 3b, open-label, non-inferiority, randomised trial. *Lancet*, **385**, 2077–2087.
- Hansson, E.C., Shams Hakimi, C., Astrom-Olsson, K., Hesse, C., Wallen, H., Dellborg, M., Albertsson, P. & Jeppsson, A. (2014) Effects of *ex vivo* platelet supplementation on platelet aggregability in blood samples from patients treated with acetylsalicylic acid, clopidogrel, or ticagrelor. *British Journal of Anaesthesia*, **112**, 570–575.
- Hastings, S., Myles, P. & McIlroy, D. (2015) Aspirin and coronary artery surgery: a systematic review and meta-analysis. *British Journal of Anaesthesia*, **115**, 376–385.
- Healey, J.S., Eikelboom, J., Douketis, J., Wallentin, L., Oldgren, J., Yang, S., Themeles, E., Heidbuchel, H., Avezum, A., Reilly, P., Connolly, S.J., Yusuf, S. & Ezekowitz, M. (2012) Perioperative bleeding and thromboembolic events with dabigatran compared with warfarin: results from the Randomized Evaluation of Long-Term Anticoagulation Therapy (RE-LY) randomized trial. *Circulation*, **126**, 343–348.
- Heidbuchel, H., Verhamme, P., Alings, M., Antz, M., Diener, H.C., Hacke, W., Oldgren, J., Sinnaeve, P., Camm, A.J. & Kirchhof, P. (2015) Updated European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist anticoagulants in patients with non-valvular atrial fibrillation. *Europace*, **17**, 1467–1507.
- Held, C., Asenblad, N., Bassand, J.P., Becker, R.C., Cannon, C.P., Claeys, M.J., Harrington, R.A., Horowitz, J., Husted, S., James, S.K., Mahaffey, K.W., Nicolau, J.C., Scirica, B.M., Storey, R.F., Vintila, M., Ycas, J. & Wallentin, L. (2011) Ticagrelor versus clopidogrel in patients with acute coronary syndromes undergoing coronary artery bypass surgery: results from the PLATO (Platelet Inhibition and Patient Outcomes) trial. *Journal*

- of the American College of Cardiology, **57**, 672–684.
- Horlocker, T.T., Wedel, D.J., Rowlingson, J.C., Enneking, F.K., Kopp, S.L., Benzon, H.T., Brown, D.L., Heit, J.A., Mulroy, M.F., Rosenquist, R.W., Tryba, M. & Yuan, C.S. (2010) Regional Anesthesia in the patient receiving antithrombotic or thrombolytic therapy; American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines (Third Edition). *Regional Anesthesia and Pain Medicine*, **35**, 64–101.
- Jamula, E., Anderson, J. & Douketis, J.D. (2009) Safety of continuing warfarin therapy during cataract surgery: a systematic review and meta-analysis. *Thrombosis Research*, **124**, 292–299.
- Kaatz, S., Douketis, J.D., White, R.H. & Zhou, H. (2011) Can the CHADS₂ score predict postoperative stroke risk in patients with chronic atrial fibrillation who are having elective non-cardiac surgery? *Journal of Thrombosis and Haemostasis*, **9**, 635.
- Kasivisvanathan, R., Abbassi-Ghadi, N., Kumar, S., Mackenzie, H., Thompson, K., James, K. & Mallett, S.V. (2014) Risk of bleeding and adverse outcomes predicted by thromboelastography platelet mapping in patients taking clopidogrel within 7 days of non-cardiac surgery. *British Journal of Surgery*, **101**, 1383–1390.
- Kearon, C. & Akl, E.A. (2014) Duration of anticoagulant therapy for deep vein thrombosis and pulmonary embolism. *Blood*, **123**, 1794–1801.
- Kearon, C. & Hirsh, J. (1997) Management of anticoagulation before and after elective surgery. *New England Journal of Medicine*, **336**, 1506–1511.
- Keeling, D., Baglin, T., Tait, C., Watson, H., Perry, D., Baglin, C., Kitchen, S. & Makris, M. (2011) Guidelines on oral anticoagulation with warfarin – fourth edition. *British Journal of Haematology*, **154**, 311–324.
- Kitchen, S., Gray, E., Mackie, I., Baglin, T. & Makris, M. (2014) Measurement of non-coumarin anticoagulants and their effects on tests of Haemostasis: Guidance from the British Committee for Standards in Haematology. *British Journal of Haematology*, **166**, 830–841.
- Korte, W., Cattaneo, M., Chassot, P.G., Eichinger, S., von Heymann, C., Hofmann, N., Rickli, H., Spannagl, M., Ziegler, B., Verheugt, F. & Huber, K. (2011) Peri-operative management of antiplatelet therapy in patients with coronary artery disease: joint position paper by members of the working group on Perioperative Haemostasis of the Society on Thrombosis and Haemostasis Research (GTH), the working group on Perioperative Coagulation of the Austrian Society for Anesthesiology, Resuscitation and Intensive Care (OGARI) and the Working Group Thrombosis of the European Society for Cardiology (ESC). *Thrombosis & Haemostasis*, **105**, 743–749.
- Kunadian, B., Thornley, A.R., Tanos, M. & Dunning, J. (2006) Should clopidogrel be stopped prior to urgent cardiac surgery? *Interactive Cardiovascular and Thoracic Surgery*, **5**, 630–636.
- Kwak, Y.L., Kim, J.C., Choi, Y.S., Yoo, K.J., Song, Y. & Shim, J.K. (2010) Clopidogrel responsiveness regardless of the discontinuation date predicts increased blood loss and transfusion requirement after off-pump coronary artery bypass graft surgery. *Journal of the American College of Cardiology*, **56**, 1994–2002.
- Lai, A., Davidson, N., Galloway, S.W. & Thachil, J. (2014) Perioperative management of patients on new oral anticoagulants. [Erratum appears in Br J Surg. 2014 Nov; 101(12):1624]. *British Journal of Surgery*, **101**, 742–749.
- Li, C., Hirsh, J., Xie, C., Johnston, M.A. & Eikelboom, J.W. (2012) Reversal of the anti-platelet effects of aspirin and clopidogrel. *Journal of Thrombosis and Haemostasis*, **10**, 521–528.
- Luckie, M., Khattar, R.S. & Fraser, D. (2009) Non-cardiac surgery and antiplatelet therapy following coronary artery stenting. *Heart*, **95**, 1303–1308.
- Makris, M. (2014) Prothrombin complex concentrate for non-vitamin K oral anticoagulant reversal: good enough for now? *Journal of Thrombosis and Haemostasis*, **12**, 1425–1427.
- Mantz, J., Samama, C.M., Tubach, F., Devereaux, P.J., Collet, J.P., Albaladejo, P., Cholley, B., Nizard, R., Barre, J., Piriou, V., Poirier, N., Mignon, A., Schlumberger, S., Longrois, D., Aubrun, F., Farese, M.E., Ravaut, P. & Steg, P.G.; Stratagem Study Group. (2011) Impact of preoperative maintenance or interruption of aspirin on thrombotic and bleeding events after elective non-cardiac surgery: the multicentre, randomized, blinded, placebo-controlled, STRATAGEM trial. *British Journal of Anaesthesia*, **107**, 899–910.
- McIlroy, D.R., Myles, P.S., Phillips, L.E. & Smith, J.A. (2009) Antifibrinolytics in cardiac surgical patients receiving aspirin: a systematic review and meta-analysis. *British Journal of Anaesthesia*, **102**, 168–178.
- Mehran, R., Baber, U., Steg, P.G., Ariti, C., Weisz, G., Witzensichler, B., Henry, T.D., Kini, A.S., Stuckey, T., Cohen, D.J., Berger, P.B., Iakovou, I., Dangas, G., Waksman, R., Antoniucci, D., Sartori, S., Krucoff, M.W., Hermiller, J.B., Shawl, F., Gibson, C.M., Chieffo, A., Alu, M., Moliterno, D.J., Colombo, A. & Pocock, S. (2013) Cessation of dual antiplatelet treatment and cardiac events after percutaneous coronary intervention (PARIS): 2 year results from a prospective observational study. *Lancet*, **382**, 1714–1722.
- Myles, P.S., Smith, J.A., Forbes, A., Silbert, B., Jayarajah, M., Painter, T., Cooper, D.J., Marasco, S., McNeil, J., Bussières, J.S. & Wallace, S. (2016) Stopping vs. Continuing Aspirin before Coronary Artery Surgery. *New England Journal of Medicine*, **374**, 728–737.
- Narouze, S., Benzon, H.T., Provenzano, D.A., Buvaendran, A., De Andres, J., Deer, T.R., Rauck, R. & Huntoon, M.A. (2015) Interventional spine and pain procedures in patients on antiplatelet and anticoagulant medications: guidelines from the American Society of Regional Anesthesia and Pain Medicine, the European Society of Regional Anaesthesia and Pain Therapy, the American Academy of Pain Medicine, the International Neuromodulation Society, the North American Neuromodulation Society, and the World Institute of Pain. *Regional Anesthesia and Pain Medicine*, **40**, 182–212.
- Nuttall, G.A., Brown, M.J., Stombaugh, J.W., Michon, P.B., Hathaway, M.F., Lindeen, K.C., Hanson, A.C., Schroeder, D.R., Oliver, W.C., Holmes, D.R. & Riha, C.S. (2008) Time and cardiac risk of surgery after bare-metal stent percutaneous coronary intervention. *Anesthesiology*, **109**, 588–595.
- Oscarsson, A., Gupta, A., Fredrikson, M., Jarhult, J., Nystrom, M., Pettersson, E., Darvish, B., Krook, H., Swahn, E. & Eintrei, C. (2010) To continue or discontinue aspirin in the perioperative period: a randomized, controlled clinical trial. *British Journal of Anaesthesia*, **104**, 305–312.
- Perry, D.J., Noakes, T.J. & Helliwell, P.S. (2007) Guidelines for the management of patients on oral anticoagulants requiring dental surgery. *British Dental Journal*, **203**, 389–393.
- Pollack, C.V. Jr, Reilly, P.A., Eikelboom, J., Glund, S., Verhamme, P., Bernstein, R.A., Dubiel, R., Huisman, M.V., Hylek, E.M., Kamphuisen, P.W., Kreuzer, J., Levy, J.H., Sellke, F.W., Stangier, J., Steiner, T., Wang, B., Kam, C.W. & Weitz, J.I. (2015) Idarucizumab for Dabigatran Reversal. *New England Journal of Medicine*, **373**, 511–520.
- Ranucci, M., Baryshnikova, E., Soro, G., Ballotta, A., De Benedetti, D. & Conti, D. (2011) Multiple electrode whole-blood aggregometry and bleeding in cardiac surgery patients receiving thienopyridines. *Annals of Thoracic Surgery*, **91**, 123–129.
- Refaai, M.A., Goldstein, J.N., Milling, T.J. Jr, Lewis, B., Goldberg-Alberts, R., Hug, B.A. & Sarode, R. (2013) Randomized phase IIb study of rapid vitamin K antagonist reversal in patients requiring an urgent surgical procedure: Four-factor prothrombin complex concentrate is superior to plasma. *Blood (ASH Annual Meeting Abstracts)*, **122**, 3588.
- Rossini, R., Musumeci, G., Visconti, L.O., Bramucci, E., Castiglioni, B., De Servi, S., Lettieri, C., Lettino, M., Piccaluga, E., Savonitto, S., Trabattini, D., Capodanno, D., Buffoli, F., Parolari, A., Dionigi, G., Boni, L., Biglioli, F., Valdatta, L., Droghetti, A., Bozzani, A., Setacci, C., Ravelli, P., Crescini, C., Staurengi, G., Scarone, P., Francetti, L., D'Angelo, F., Gadda, F., Comel, A., Salvi, L., Lorini, L., Antonelli, M., Bovenzi, F., Cremonesi, A., Angiolillo, D.J. & Guagliumi, G. (2014) Perioperative management of antiplatelet therapy in patients with coronary stents undergoing cardiac and non-cardiac surgery: a consensus document from Italian cardiological, surgical and anaesthesiological societies. *EuroIntervention*, **10**, 38–46.
- Rossini, R., Musumeci, G., Capodanno, D., Lettieri, C., Limbruno, U., Tarantini, G., Russo, N.,

- Calabria, P., Romano, M., Inashvili, A., Sirbu, V., Guagliumi, G., Valsecchi, O., Senni, M., Gavazzi, A. & Angiolillo, D.J. (2015) Perioperative management of oral antiplatelet therapy and clinical outcomes in coronary stent patients undergoing surgery: results of a multicentre registry. *Thrombosis and Haemostasis*, **113**, 272–282.
- Savonitto, S., Caracciolo, M., Cattaneo, M. & De Servi, S. (2011) Management of patients with recently implanted coronary stents on dual antiplatelet therapy who need to undergo major surgery. *Journal of Thrombosis & Haemostasis*, **9**, 2133–2142.
- Schulman, S., Hwang, H.G., Eikelboom, J.W., Kearon, C., Pai, M. & Delaney, J. (2014) Loading dose vs. maintenance dose of warfarin for reinitiation after invasive procedures: a randomized trial. *Journal of Thrombosis and Haemostasis*, **12**, 1254–1259.
- Schulman, S., Carrier, M., Lee, A.Y., Shivakumar, S., Blostein, M., Spencer, F.A., Solymoss, S., Barty, R., Wang, G., Hedde, N. & Douketis, J.D.; Perioperative Management of Dabigatran Study Group. (2015) Perioperative Management of Dabigatran: A Prospective Cohort Study. *Circulation*, **132**, 167–173.
- Shakur, H., Roberts, I., Bautista, R., Caballero, J., Coats, T., Dewan, Y., El-Sayed, H., Gogichaishvili, T., Gupta, S., Herrera, J., Hunt, B., Iribhogbe, P., Izurieta, M., Khamis, H., Komolafe, E., Marrero, M.A., Mejia-Mantilla, J., Miranda, J., Morales, C., Olaomi, O., Oildashi, F., Perel, P., Peto, R., Ramana, P.V., Ravi, R.R. & Yuthakasemsunt, S. (2010) Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet*, **376**, 23–32.
- Sherwood, M.W., Douketis, J.D., Patel, M.R., Piccini, J.P., Hellkamp, A.S., Lokhnygina, Y., Spyropoulos, A.C., Hankey, G.J., Singer, D.E., Nessel, C.C., Mahaffey, K.W., Fox, K.A., Califf, R.M. & Becker, R.C. (2014) Outcomes of temporary interruption of rivaroxaban compared with warfarin in patients with nonvalvular atrial fibrillation: results from the rivaroxaban once daily, oral, direct factor Xa inhibition compared with vitamin K antagonism for prevention of stroke and embolism trial in atrial fibrillation (ROCKET AF). *Circulation*, **129**, 1850–1859.
- Shi, J., Ji, H., Ren, F., Wang, G., Xu, M., Xue, Y., Chen, M., Qi, J. & Li, L. (2013) Protective effects of tranexamic acid on clopidogrel before coronary artery bypass grafting: a multicenter randomized trial. *JAMA Surgery*, **148**, 538–547.
- Siegal, D., Yudin, J., Kaatz, S., Douketis, J.D., Lim, W. & Spyropoulos, A.C. (2012) Periprocedural heparin bridging in patients receiving vitamin K antagonists: systematic review and meta-analysis of bleeding and thromboembolic rates. *Circulation*, **126**, 1630–1639.
- Siegal, D.M., Curnutte, J.T., Connolly, S.J., Lu, G., Conley, P.B., Wiens, B.L., Mathur, V.S., Castillo, J., Bronson, M.D., Leeds, J.M., Mar, F.A., Gold, A. & Crowther, M.A. (2015) Andexanet Alfa for the Reversal of Factor Xa Inhibitor Activity. *New England Journal of Medicine*, **373**, 2413–2424.
- Singla, S., Sachdeva, R. & Uretsky, B.F. (2012) The risk of adverse cardiac and bleeding events following noncardiac surgery relative to antiplatelet therapy in patients with prior percutaneous coronary intervention. *Journal of the American College of Cardiology*, **60**, 2005–2016.
- Smith, P.K., Goodnough, L.T., Levy, J.H., Poston, R.S., Short, M.A., Weerakkody, G.J. & Lenarz, L.A. (2012) Mortality benefit with prasugrel in the TRITON-TIMI 38 coronary artery bypass grafting cohort: risk-adjusted retrospective data analysis. *Journal of the American College of Cardiology*, **60**, 388–396.
- Soo, C.G., Torre, P.K., Yolland, T.J. & Shatwell, M.A. (2016) Clopidogrel and hip fractures, is it safe? A systematic review and meta-analysis. *BMC Musculoskeletal Disorders*, **17**, 136.
- Sorensen, R., Hansen, M.L., Abildstrom, S.Z., Hvelplund, A., Andersson, C., Jorgensen, C., Madsen, J.K., Hansen, P.R., Kober, L., Torp-Pedersen, C. & Gislason, G.H. (2009) Risk of bleeding in patients with acute myocardial infarction treated with different combinations of aspirin, clopidogrel, and vitamin K antagonists in Denmark: a retrospective analysis of nationwide registry data. *Lancet*, **374**, 1967–1974.
- Sousa-Uva, M., Storey, R., Huber, K., Falk, V., Leite-Moreira, A.F., Amour, J., Al-Attar, N., Ascione, R., Taggart, D. & Collet, J.P. (2014) Expert position paper on the management of antiplatelet therapy in patients undergoing coronary artery bypass graft surgery. *European Heart Journal*, **35**, 1510–1514.
- Spyropoulos, A.C. & Douketis, J.D. (2012) How I treat anticoagulated patients undergoing an elective procedure or surgery. *Blood*, **120**, 2954–2962.
- Steib, A., Barre, J., Mertes, M., Morel, M.H., Nathan, N., Ozier, Y., Treger, M. & Samama, C.M. (2010) Can oral vitamin K before elective surgery substitute for preoperative heparin bridging in patients on vitamin K antagonists? *Journal of Thrombosis & Haemostasis*, **8**, 499–503.
- Thiele, T., Sumnig, A., Hron, G., Muller, C., Althaus, K., Schroeder, H.W. & Greinacher, A. (2012) Platelet transfusion for reversal of dual antiplatelet therapy in patients requiring urgent surgery: a pilot study. *Journal of Thrombosis and Haemostasis*, **10**, 968–971.
- Veitch, A.M., Vanbiervliet, G., Gershlick, A.H., Boustiere, C., Baglin, T.P., Smith, L.A., Radaelli, F., Knight, E., Gralnek, I.M., Hassan, C. & Dumonceau, J.M. (2016) Endoscopy in patients on antiplatelet or anticoagulant therapy, including direct oral anticoagulants: British Society of Gastroenterology (BSG) and European Society of Gastrointestinal Endoscopy (ESGE) guidelines. *Gut*, **65**, 374–389.
- Vilahr, G., Choi, B.G., Zafar, M.U., Viles-Gonzalez, J.F., Vorchheimer, D.A., Fuster, V. & Badimon, J.J. (2007) Normalization of platelet reactivity in clopidogrel-treated subjects. *Journal of Thrombosis & Haemostasis*, **5**, 82–90.
- Ward, C., Conner, G., Donnan, G., Gallus, A. & McRae, S. (2013) Practical management of patients on apixaban: a consensus guide. *Thrombosis Journal*, **11**, 27.